



Purina Mills



# Powerful Performance. Winning Results.



## PRELUDE™ 409

Feed Honor® Show Chow® PRELUDE™ 409 to 25 to 50 lb pigs. This feed can be fed to heavier pigs.

PRELUDE™ 409 contains Safe-Guard® dewormer. PRELUDE™ can also be used to deworm heavier growing/finishing showpigs. Feed 25 lbs of PRELUDE™ 409 for every 100 lbs of bodyweight.

Honor® Show Chow® PRELUDE™ 409 is supported by Purina Mills® Animal Nutrition. Get your newly weaned pigs on the PRELUDE™ program today to Unleash Your Genetic Investment.

FEATURES	BENEFITS
Formulated with very high quality animal protein sources	Very palatable, newly weaned pigs get on feed fast and look great
High levels of protein, essential amino acids (1.4% lysine), and crude fat (7.0%)	Nutrient dense to support intake and growth of young pigs
Organic acids, yucca schdigera extract, and high levels of vitamins.	Acidified to enhance digestibility, yucca to promote intake and air quality, and high in fat and water soluble vitamins for small pigs
3/32" pellets	Extremely small pellet for improved intake
Two medication options (CTC/DEN & L200 & FBZ36)	Prevent and control respiratory and digestive tract diseases. Lincomycin and Safe-Guard® – protection against PPLO (mycoplasma pneumonia) & feed through dewormer
Proprietary blend of ingredients	Produces full middles, muscle shape and bloom
Moderately high lactose levels	Easy transition from PRELUDE™ 209 to PRELUDE™ 409

Visit our website to download this product form.



[www.HonorShowChow.com](http://www.HonorShowChow.com)

Regional Availability

Check your current price list for availability in your region, or contact your local Purina Dealer or Representative for more information.

For more information on this product, visit our website at [www.HonorShowChow.com](http://www.HonorShowChow.com) or call Purina Customer Service at 1-800-227-8941

# PRELUDE™ 409 L200/FBZ36

## GUARANTEED ANALYSIS

Crude Protein, %	(min)	20.0
Lysine, %	(min)	1.40
Crude Fat, %	(min)	7.00
Crude Fiber, %	(max)	3.50
Calcium, % (Ca)	(min)	0.60
Calcium, % (Ca)	(max)	1.10
Phosphorus, % (P)	(min)	0.60
Salt, % (Na Cl)	(min)	0.60
Salt, % (Na Cl)	(max)	1.10
Selenium, ppm (Se)	(min)	0.30
Zinc, ppm (Zn)	(min)	500

**Complete Swine Starter**  
**Medicated for pigs weighing 25-50 lbs.**

For reduction in the severity of swine mycoplasmal pneumonia caused by mycoplasma hypopneumoniae. Feed as the sole ration for 21 days. For the removal of : Lungworms: Metastrongylus apri, Metastrongylus pudendotectus. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (Ascaris sum); nodeular (Oesophafostomum dentatum, Ornithostrongylus quadrispinulaturm); small stomach (Hyostromylus rubidus); Adult and larvae (L2,L3,L4 stages intestinal mucosal forms) whipworm (Trichuris suis); Kidneyworm: Audlt and larvae Stephanurus dentatus.

**Active Drug Ingredient:** Lincomycin 200g/ton  
Fenbendazole 36g/ton

## FEEDING DIRECTIONS

Feed HSC PRELUDE™ 409 L200/FBZ36 as the sole ration in complete feed for a period of from 3-12 days so to provide a total intake of 9 mg/kg of bodyweight of Fenbendazole.

**Warning:**

Do not slaughter swine for human consumption for 6 days following last treatment when fed at 100 or 200 g/ton of complete feed.

**Caution:**

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not to be fed to swine that weigh more than 250 lb.